

Strong Jump Traceability I

Rod Downey
Victoria University
Wellington
New Zealand

BACKGROUND REFERENCES

- ▶ Calibrating Randomness, (Downey, Hirschfeldt, Nies, Terwijn) Bulletin Symbolic Logic, 2006
- ▶ Algorithmic Randomness and Complexity, (Downey and Hirschfeldt) Springer-Verlag
- ▶ Randomness and Computability (Nies) OUP

REFERENCES

- ▶ Strong Jump-Traceability I : the Computably Enumerable Case, (Cholak, Downey, Greenberg) *Advances in Math.*
- ▶ Strong jump-traceability II: (Downey and Greenberg) *Israel J Math.*
- ▶ Lowness properties and approximations of the jump. (Figueira, Nies, Stephan), *APAL*
- ▶ *K*-trivial degrees and the jump-traceability hierarchy, Barmpalias, Downey and Greenberg. *Proc AMS*
- ▶ Beyond strong jump traceability, (Ng Keng Meng) *Proc LMS*

- ▶ Characterising the strongly jump-traceable sets via randomness, Greenberg, Hirschfeldt and Nies, Adv Math.
- ▶ Strong jump-traceability and Demuth randomness, Greenberg and Turetsky. to appear Proc. LMS
- ▶ Benign cost functions and lowness notions, Greenberg and Nies. JSL.
- ▶ Pseudo-jump operators and SJTHard sets, (Downey and Greenberg) to appear Advances in Math.
- ▶ Inherent enumerability of strong jump traceability, (Diamondstone, Greenberg, Turetsky)

NOTATION

- ▶ Real is a member of Cantor space 2^ω with topology with basic clopen sets $[\sigma] = \{\sigma\alpha : \alpha \in 2^\omega\}$ whose measure is $\mu([\sigma]) = 2^{-|\sigma|}$.
- ▶ $\alpha \upharpoonright n$ is the first n bits of α .
- ▶ Strings = members of $2^{<\omega} = \{0, 1\}^*$.

KOLMOGOROV COMPLEXITY

- ▶ Capture the incompressibility paradigm. Random means hard to describe, incompressible: e.g. 1010101010.... (10000 times) would have a short program.
- ▶ A string σ is random iff the only way to describe it is by hardwiring it. (Formalizing the Berry paradox)
- ▶ Want the **bits** of τ to describe σ if $U(\tau) = \sigma$ for a device (Turing machine) U . Write $C(\sigma) = |\tau|$ for the **shortest** such τ , and can use a universal machine.
- ▶ Plain complexity like this has τ providing itself and its length so this is circumvented by using prefix-free complexity (telephone numbers) giving K for prefix-free machines.
- ▶ Using this, Levin, Chaitin, Schnorr proved that there are reals with $K(\alpha \upharpoonright n) \geq^+ n$ for all n , called **1-random**, and coinciding with earlier ones avoiding all effective null sets.

MARTIN-LÖF RANDOMNESS

- ▶ Generalized effective statistical test: an **effectively shrinking** computable collection of open sets: $\{U_n \mid n \in \mathbb{N}\}$ with $\mu(U_n) \leq 2^{-n}$.
- ▶ e.g. every second bit is 0: $U_1 = \{[10], [00]\}$, etc.
- ▶ For all such **Martin-Löf** tests $A \notin \bigcap_n U_n$.
- ▶ Other notions: **Schnorr randomness** ($\mu(U_n) = 2^{-n}$), **Demuth Randomness** (ω -effective approximations and other acceptances criterion), etc.

- ▶ As with life, relationships here are complex (Solovay)

$$K(x) = C(x) + C^{(2)}(x) + \mathcal{O}(C^{(3)}(x)).$$

and

$$C(x) = K(x) - K^{(2)}(x) + \mathcal{O}(K^{(3)}(x)).$$

- ▶ These 3's are **sharp** (Solovay) That is, for example, $K = C + C^2 + C^3 + \mathcal{O}(C^4)$ is NOT true.

LOWNESS

- ▶ I would like to discuss the remarkable story of lowness generating *K*-triviality and then sjt.
- ▶ First sjt was an apparent artifact
- ▶ Then proved to be intimately related with randomness
- ▶ Later (also) giving insight into computability itself.
- ▶ I will try to explain the **little boxes** method, which is new and poorly understood.
- ▶ Theme: to what extent do **computational lowness** (the extent to which sets resemble computable ones) and being **far from random** align themselves?

KEY FACTS

▶ THEOREM (CHAITIN)

There is a constant d such that for all c , and n ,

$$|\{\nu : |\nu| = n \wedge C(\nu) \leq C(n) + c\}| \leq d2^c.$$

THEOREM (LEVIN, CHAITIN)

There is a constant d such that for all c and n ,

$$|\{\nu : |\nu| = n \wedge C(\nu) \leq C(n) + c\}| \leq d2^c.$$

INFORMATION CHARACTERIZATION OF COMPUTABILITY

- ▶ Chaitin proved that a real A is computable iff for all n , $C(A \upharpoonright n) \leq^+ \log n$, iff $C(A \upharpoonright n) \leq^+ C(n)$.
- ▶ This is proven using the fact that a Π_1^0 class with a finite number of paths has only computable paths, combined with the Counting Theorem
 $\{\sigma : C(\sigma) \leq C(n) + c \wedge |\sigma| = n\} \leq d2^c$. (Using the Meyer-Loveland Technique below)
- ▶ Meyer(-Loveland) had earlier shown A is computable iff $C(A \upharpoonright n|n) \leq c$ for some c and all n .

THE MEYER-LOVELAND TECHNIQUE

- ▶ If $C(\alpha \upharpoonright n|n) \leq c$ then there are only c programmes possibly computing initial segments of α .
- ▶ This computes a tree of *strings* of maximal width c .
- ▶ Therefore only at most c paths. Say \hat{c} .
- ▶ Imagine the situation that there is only *one* path in a tree of maximal width 2.
- ▶ Enumerate until only one remains.

K-TRIVIALITY

- ▶ What is a consequence of $K(A \upharpoonright n) \leq^+ K(n)$ for all n ? We call such reals **K-trivial**. Does A K-trivial imply A computable?
- ▶ Write $A \in KT(d)$ iff for all n , $K(A \upharpoonright n) \leq K(n) + d$.

THE ARGUMENT FAILS

- ▶ It is still true that $\{\sigma : K(\sigma) \leq K(|\sigma|) + d\}$ is $O(2^d)$, so it would appear that we could run the Π_1^0 class argument used for C . But no...
- ▶ The **problem** is that we don't know $K(n)$ in any computable interval, therefore the tree of K -trivials we would construct would be a Π_1^0 class **relative to \emptyset'** .

THEOREM (CHAITIN, ZAMBELLA)

There are only $O(2^d)$ members of $KT(d)$. They are all Δ_2^0 .

THEOREM (SOLOVAY)

*There are noncomputable *K*-trivial reals.*

THEOREM (ZAMBELLA)

Such reals can be c.e. sets.

A REMARKABLE CLASS

- ▶ *K*-trivials form a remarkable class as we will see.
- ▶ First they solve Post's problem.
- ▶ Theorem: (DHNS) If *A* is *K*-trivial then $A <_T \emptyset'$.

Similar methods allow for us to show the following

THEOREM (NIES)

All K -trivials are superlow $A' \equiv_{tt} \emptyset'$, and are tt -bounded by c.e. K -trivials. In fact they are **Jump Traceable** as we see below.

Thus triviality is essentially an “enumerable” phenomenon.

There are other antirandomness notions.

DEFINITION (KUČERA AND TERWIJN)

We say A is low for randomness iff the reals Martin-Löf random relative to A are exactly the Martin-Löf random reals.

DEFINITION (HIRSCHFELDT, NIES, STEPHAN)

A is a base for randomness iff $A \leq_T B$ with B A -random.

THEOREM

The following are equivalent to being K-trivial.

- (I) (Nies) *A is low for randomness.*
- (II) (Hirschfeldt and Nies) *A is K-low in that $K^A =^+ K$.*
- (III) (Hirschfeldt, Nies, Stephan) *A is a base for randomness.*
- (IV) (Downey, Nies, Weber, Yu+Nies, Miller) *A is low for weak-2-randomness.*
- (V) *+ 15 others!*

QUESTIONS AND A PROPER SUBCLASS

It is open if this is the same as a number of other “cost function” classes such as the reals which are Martin-Löf coverable. It is known there is a proper subclass defined by cost function.

DEFINITION

- ▶ (Zambella, Terwijn, later Nies) Let h be an order. We say that A is **jump traceable** for the order h iff there is a computable collection of c.e. sets $W_{g(e)}$ with $|W_{g(e)}| < h(e)$ and $J^A(e) \in W_{g(e)}$, for every partial A -computable function J^A .
- ▶ (Figueira, Nies, Stephan) A is **strongly jump traceable** iff it is jump traceable for **every** computable order.

Think about classical set theory notions of capturing a function by specifying possibilities. (Raisonnier, Shelah, ..., Zambella).

COMBINATORIAL IDEAS

Inspired by the result

THEOREM (TERWIJN-ZAMBELLA, THEN BEDREGAL, KJOS-HANSEN, NIES, STEPHAN)

A is low for Schnorr randomness iff A is computably traceable. That is, for every function $f \leq_T A$ there is a canonical collection of finite sets $\{D_{g(n)} \mid n \in \mathbb{N}\}$ such that $|D_{g(n)}| \leq n + 1$ and for all n , $f(n) \in D_{g(n)}$.

This generalized old tracing notions such as highness (Martin) lowness (Soare), hyperhyperimmunity (Miller-Martin).

THEOREM (NIES)

A is K-triv implies that there is an order $h (n \log n)$ relative to which A is jump traceable.

Improved to $M \log n$ for **some** M by Hölzl, Kräling, Merkle.

THEOREM (FIGUEIRA, NIES, STEPHAN)

Noncomputable sjt c.e. sets exist.

THEOREM (FIGUEIRA, NIES, STEPHAN)

*If A is **strongly superlow** the A is sjt.*

strongly superlow means that A' has very tame approximations

THEOREM (FIGUEIRA, NIES, STEPHAN)

Sjt is equivalent to $C(n) \leq^+ C^A(n) + h(C^A(n))$ for all orders h .

ie A is **lowly for C**

THEOREM (CHOLAK, DOWNEY, GREENBERG)

The c.e. sjt's are a *proper* subclass of the K -trivials. They form an ideal.

Turetsky has recently shown that there is a K -trivial which is not $o(\log n)$ -jump traceable.

THEOREM (DOWNEY, GREENBERG)

If A is sjt then A is Δ_2^0

THEOREM (DIAMONDSTONE, GREENBERG, TURETSKY)

If A is sjt then $A \leq_T B$ with B sjt and c.e.

COROLLARY (DIAMONDSTONE, GREENBERG, TURETSKY)

A is sjt is equivalent to A is strongly super low.

RANDOMNESS

- ▶ It might seem that sjt's are an *artifact* of randomness and potential combinatorial characterizations of notions. However (Noam will be discussing these results):

THEOREM (GREENBERG, HIRSCHFELDT, NIES)

sjt = $\text{superlow}^\diamond = \text{superhigh}^\diamond$. Here C^\diamond is exactly the c.e. sets below all random members of C .

THEOREM (GREENBERG, NIES)

A is *sjt* iff it is computable from all Δ_2 MRL which are not weakly Demuth random.

THEOREM (KUCERA-NIES, GREENBERG-TURETSKY)

A c.e. degree **a** is *sjt* iff it is computable from some **Demuth** random real.

- ▶ Roughly need orders $\sqrt{\log n}$, $o(\log n)$. Is there a combinatorial characterization?
- ▶ Conjecture : A is K -trivial iff A is jump traceable for all computable orders h with $\sum_{n \geq 1} 2^{-h(n)} < \infty$.

A HINT OF THE PROOF TECHNIQUES

- ▶ To show that if A and B are c.e. sjt, so is $A \oplus B$.
- ▶ Given h we can construct a slower order k such that if A and B are jump traceable via k then $A \oplus B$ is jump traceable via h
- ▶ Opponent gives: $W_{p(x)}$ jump tracing A and $W_{q(x)}$ jump tracing B , such that $|W_{p(x)}|, |W_{q(x)}| < k(x)$.
- ▶ We: V_z tracing $J^{A \oplus B}(z)$ with $|V_z| < h(z)$.

TWO OBSTACLES

- ▶ We see an **apparent** jump computation $J^{A \oplus B}(x) \downarrow [s]$.
- ▶ Should we believe? We only have $h(x)$ many slots in the trace V_x to put possible values.
- ▶ Opponent can change A or B after stage s on the use.
- ▶ We build parts of jump (recursion thm) testing A and B
- ▶ Basic idea: For some $a = a(x)$ and $b = b(x)$ we will define

$$J^B[s](b) = j_B(x, s) \text{ and } J^A[s](a) = j_A(x, s),$$

where $j_C(x, s)$ denotes the C -use of the $J^{A \oplus B}(x)[s]$ computation.

- ▶ Ignore *noncompletion*: that is the $A \oplus B$ computation changes **before** these procedures return.
- ▶ Simplest case: $W_{p(a)}$ and $W_{q(b)}$ were of size 1 (1-boxes)
- ▶ Then if return: $A \oplus B$ **is** correct
- ▶ Now 2-boxes. If the $A \oplus B$ computation is wrong, at least one of the A or B ones are too.

- ▶ If we are lucky and there is are false jump computations in both of the $W_{p(a)}$ and $W_{q(b)}$.
- ▶ The 2-boxes are now, in effect 1-boxes. (Very good)
- ▶ Can't allow to only point at one side. Use up all the 2-boxes.
- ▶ For example if always the A sides was the wrong part, and there were k 2-boxes then after k attacks, all the 2-boxes would be useless and the information in the B -side is correct, hence the box is used.

MULTIPLE BOXES

- ▶ Idea: use multiple 2-boxes. E.g. at the beginning use two 2-boxes for the **same** computation.
- ▶ A side was wrong. Then now we have *two* **promoted** 1-boxes.
- ▶ Since the *A*-computation now must be correct, if the believed computation is wrong, it must be the *B* side which wrong the next time, now creating a new *B*-1-box. Finally the third time we test, we would have two 1-boxes.

NON-RETURN

- ▶ Now we face the ignored problem. We test and **before** the computation returns, the jump computation is changed by an *A* or *B* change, but **possibly** one of the *A* or *B* uses **is** correct. Now **nothing** is promoted. This seems very bad.
- ▶ Even with 1-boxes.
- ▶ Use descending sequences of boxes, and monster boxes.
- ▶ Complicated, combinatorial.

- ▶ The idea is that for a computation whose target is, say, 2-boxes, begin ever further out. Begin by testing at, say, s -boxes.
- ▶ Monster boxes called **metaboxes**.
- ▶ If **both** A and B return at the s -box, go to $s - 1$ etc. Only believe if you get back to the 2-boxes. The idea that a failure at k **promotes** $k + 1, \dots, s$ -boxes, at least on one side.
- ▶ A combinatorial argument if used to show that cannot favour one side forever.

PROPER SUBCLASS OF THE *K*-TRIVIALS

- ▶ How to make a properly ω -c.e. *K*-trivial?
- ▶ Use **descending costs**....
- ▶ If the trace grows slowly enough then can make *K*-trivial and **not** jt at that order. Much the same idea, the key point being the to change the trace and use a a box location, the **use** is very big, and the opponent needs more tailweight.

NG KENG MENG'S THEOREMS

- ▶ The c.e. sjt's are Π_4^0 complete.
- ▶ This solves a problem of Nies: there is no minimal order.

BEYOND JUMP TRACEABILITY

- ▶ Say A is **C-sjt** iff for all orders h^B , for $B \in C$, A is h^B -jt.
- ▶ (Ng) No real is C-sjt where $C = \Delta_2$.
- ▶ (Ng) There are c.e. reals sjt for all c.e. sets.
- ▶ (Ng) They cannot be promptly simple, the first such class.
- ▶ (Ng) **No** real is K^B -trivial for all B , or c.e. B .